

# Suggested Three Year Worming Cycle

(by Nick Bowen BVetMed, CertEP, MRCVS, Chalgrave Equine Clinic near Dunstable)

## Year One

(Main wormer is from group A i.e. a macrocyclic lactone)

| Jan | Feb | Mar    | Apr                                                                          | May | Jun    | Jul | Aug | Sept   | Oct                                                                         | Nov | Dec                                                                   |
|-----|-----|--------|------------------------------------------------------------------------------|-----|--------|-----|-----|--------|-----------------------------------------------------------------------------|-----|-----------------------------------------------------------------------|
|     |     | Equest | Strongid-P<br>or Pyratape P<br>(Double dose<br>for tapeworm)<br>or Equitape. |     | Equest |     |     | Equest | Strongid-P<br>or Pyratape P<br>(Double dose<br>for tapeworm<br>or Equitape) |     | Equest<br>(bots & some<br>of the encysted<br>small redworm<br>larvae) |

## Year Two

(Main wormer is from group B i.e. a pyrimidine)

| Jan                         | Feb | Mar | Apr                                                          | May | Jun                                                                                                                                                                                | Jul | Aug | Sept | Oct                                                          | Nov                                                           | Dec               |
|-----------------------------|-----|-----|--------------------------------------------------------------|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----|------|--------------------------------------------------------------|---------------------------------------------------------------|-------------------|
| Strongid-P<br>or Pyratape P |     |     | Strongid-P<br>or Pyratape P<br>(Double dose<br>for tapeworm) |     | 6 weekly dosing with Strongid P or<br>Pyratape P (The recommended dosage<br>interval is 6 weeks. This could be<br>extended if your management is good.<br>See text in the article) |     |     |      | Strongid P<br>or Pyratape P<br>(Double dose<br>for tapeworm) | Panacur<br>Equine Guard<br>(encysted small<br>redworm larvae) | Eqvalan<br>(bots) |

## Year Three

(Main wormer is from group A i.e. a macrocyclic lactone)

| Jan                                                                                                                                                       | Feb     | Mar | Apr                      | May | Jun | Jul     | Aug | Sept | Oct                      | Nov | Dec               |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|---------|-----|--------------------------|-----|-----|---------|-----|------|--------------------------|-----|-------------------|
|                                                                                                                                                           | Eqvalan |     | Equimax<br>(dual wormer) |     |     | Eqvalan |     |      | Equimax<br>(dual wormer) |     | Eqvalan<br>(bots) |
| <p>(The recommended dosage interval is 8-10 weeks.<br/>This could be extended if your management is good.<br/>See text article on previous two pages)</p> |         |     |                          |     |     |         |     |      |                          |     |                   |



Bots



Large redworms



Small redworms



Tapeworm